

Quality Index Method developed for Frigate tuna (*Auxis thazard*)

K.W.S. ARIYAWANSA*¹, D.N. WIJENDRA² AND S.P.S.D. SENADHEERA¹

¹Post Harvest Technology Division

National Aquatic Resources Research and Development Agency
Crow Island, Colombo 15, Sri Lanka

²Department of Chemistry, University of Sri Jayawardenapura
Sri Lanka

*Corresponding author: sujeewa@nara.ac.lk

Abstract

An investigation was carried out to develop and evaluate a Quality Index Method (QIM) scheme for Frigate tuna (*Auxis thazard*) for the prediction of past and remaining storage time in ice. The quality index method (QIM) provides weighted evaluation of the key parameters in deterioration of individual species, assigning demerit points according to the importance of each parameter. For the development and evaluation of QIM scheme, Frigate tuna were stored in ice and assessed raw and cooked. The QIM developed for raw Frigate tuna comprised of 11 parameters (appearance of skin and stiffness: cornea, form and colour of eyes; colour, smell and mucus of gills; condition of viscera; colour of blood and fillets) covering attributes, which gave a total of 25 demerit points. Sensory analysis of cooked Frigate tuna using Torry scheme were carried out in parallel to determine the shelf life. In order to obtain more information about the quality of Frigate tuna, Total viable counts (TVC), hydrogen sulphide (H_2S) producing bacteria, trimethylamine (TMA), total volatile basic nitrogen (TVN) content and pH value were determined.

The sensory evaluation was carried out by a panel of 6-7 judges and the rejection level was found to be 22 days. The QIM scheme developed for Frigate tuna showed a linear relationship between QIM scores and storage time in ice, ($r^2 = 0.9378$) with slope of 0.755. The TVC varied from 10^2 cfu/g to 10^6 cfu/g within 7 to 22 days and H_2S producing bacteria at 10^2 cfu/g within 22 days of storage in ice. The TMA and TVN amounts increased with time and the amounts ranged from 1.2 mg/100g to 2.5 mg/100g and 17.2 mg/100g to 34.9 mg/100g respectively. The variation of pH ranged from 5.4 to 5.9 within the storage.

The QIM calibration curve obtained for Frigate tuna indicates its applicability to determine the storage life of fish in ice. Further, it will facilitate the requirements of buyers and sellers while fulfilling the demands of inspection authorities and the consumers.

Introduction

Improving the quality and the availability of fish for human consumption is an important area for management action. Handling and preservation of catch on board need to be improved, by using ice, chilled water, refrigeration/ freezing systems, etc. The fish that is landed must be kept clean and stored in ice, in refrigerated rooms or transported in ice, in refrigerated trucks depending on the duration of storage or transportation.

The freshness of fish deteriorates with time. Most of the changes resulting from loss of freshness can be evaluated by sight, touch, smell and taste which determine consumer acceptance. It is therefore extremely useful to have methods for describing the sensory properties of fish as a means of ascertaining their initial sensory characteristics and any changes undergone by the product in the course of storage.

The quality index method (QIM) is a very effective tool and a rapid and simple descriptive sensory method to determine freshness and quality (Bremner 1985; Bremner et al. 1985). QIM essentially evaluates those sensory parameters and attributes that change most significantly in each species during storage period (Huss 1995). By using QIM, the fish inspector can give estimation of the past and remaining storage time. In addition, the knowledge of shelf life of fish species would allow fishermen, processors and retailers better control of their markets.

Regulations in European Union require freshness grading of most fish to be marketed within the union. Further, the fish inspection department has to perform quality inspection of fish prior to export in order to issue health certificate in accordance with EU regulations (EU Directive 1991). Fisheries and Aquatic Resources Act of 1996 also describes the freshness category of selected fish species (Fish Products (Export) Regulation 1998).

The aim of this study was to develop a quality index specific for raw fish of Frigate tuna (*Auxis thazard*) and predict the shelf life during ice storage and to compare the sensory analysis with microbial counts (total viable counts and H₂S producing bacteria) and chemical measurements.

Materials and Methods

A total of 70 fresh Frigate tuna (*Auxis thazard*) caught in Matara, the south cost of Sri Lanka were used in this experiment. The fish which were

Quality index for Frigate tuna

kept in ice 18 hrs after capture were brought to NARA laboratory. The first batch of Frigate tuna (30) was used for training of assessors, development of quality index and for photography. A batch of 40 fish was stored at 0°C to 2°C in iced boxes until analysis, which was used for test measurements and sensory evaluation.

The sensory evaluation of raw and cooked Frigate tuna were carried out in parallel on each sampling day (no. 1, no. 7, no. 12, no. 16, no. 19 and no. 22 days after capture) along with measurements of microbiological, chemical and pH. Each sampling day, 5 Frigate tuna were used. Three fish were analyzed whole by QIM (Table 1). Pieces from one fish were steam cooked and evaluated with the Torry scheme (Table 2) for odour and flavour. Sensory assessments of cooked fish were carried out in parallel. Rest of the fish were used for microbiological analysis for TVC and H₂S producing bacteria (Gram et al. 1989), chemical analysis for TVN and TMA (Horwits 1980) and pH measurements. A panel of 5-7 trained judges evaluated the fish individually for each quality parameter in the scheme. Selected pictures were used to develop the QIM scheme for raw fish.

Results

The results from sensory evaluation, chemical measurements, microbial counts and pH are presented in relation to storage time of Frigate tuna in ice. The sum of scores evaluated ranging from 0 for the best quality fish to 25 for the poorest quality fish. According to the QIM scheme (Table 1) quality is presented as Quality Index (QI). The QI was linearly related to storage period ($R^2 = 0.937$) (Fig. 1).

The scores for all quality attributes of the QIM increased with storage time in ice (Figure 2). On the first day appearance of fish was scored 0-1, as the skin of fish was bright. On the 22nd day fish were scoring 2 as the appearance of skin was dull. The skin did not become very soft after 22 days of storage in ice. The scores of the quality attributes of gills increased with storage time. The colour approached maximum after 22 days in ice. The smell of gills did not reach the maximum of score of 3, the average score of which was however above 1 after 16 days. The scores for gill odour appeared to have the same trend as gill mucus, but the average score reached around 2 at the end of storage time. The scores for viscera increased with storage time in ice, constantly.

Table 1. Quality Index method (QIM) scheme for Frigate tuna (*Auxis thazard*).

Quality parameter		Description	Score
Appearance:	Skin	Bright, iridescent pigmentation	0
		Rather dull, becoming discolored	1
		Dull	2
	Stiffness	In rigor	0
		Firm, elastic	1
		Soft	2
	Very soft	3	
Eyes:	Cornea	Clear	0
		Opalescent	1
		Milky	2
	Form	Convex	0
		Flat, slightly sunken	1
		Sunken, concave	2
	Colour of Pupil	Black	0
		Opaque	1
	Grey	2	
Gills:	Colour	Bright	0
		Less colored, becoming discoloured	1
		Discolored, brown spots	2
		Brown, discolored	3
	Smell	Fresh, seaweedy, metallic	0
		Neutral, grassy, musty	1
		Yeast, bread, beer, sour milk	2
		Acetic acid, sulphuric	3
	Mucus	Clear	0
		Milky	1
		Milky, dark, opaque	2
Viscera	Whole	0	
	Beginning to get dissolved	1	
	Viscera dissolved	2	
Blood:	Color	Red	0
		Dark red	1
		Brown	2
Fillets:	Color	Translucent, bluish	0
		Waxy, milky	1
		Opaque, yellow, brown spots	2
Quality Index			Sum: (0-25)

Quality index for Frigate tuna

Table 2. The score sheet for Torry scheme of freshness for cooked Frigate tuna (*Auxis thazard*).

Score	Odour	Flavour
10	Initially weak odour of boiled cod liver, fresh oil, starchy	Boiled cod liver Watery, metallic
9	Shellfish, seaweed, Boiled meat Oil, cod liver	Oily, boiled cod liver Sweet, meaty, Characteristic
8	Loss of odour, Neutral odour	Sweet and characteristic flavours but reduced in intensity
7	Wood shavings, woodsap, vanillin	Neutral
6	Condensed milk, Boiled potato	Inspid
5	Milk jug odours, boiled clothes-like	Slight sourness, trace of "off"-flavours, rancid
4	Lactic acid, sour milk, TMA	Slight bitterness, sour, "off"-flavours, TMA, rancid
3	Lower fatty acids (e.g. acetic or butric acids) composed grass, soapy, turnipy, tallowy	Strong bitter, rubber, slight sulphide, rancid

Torry scheme

Using Torry scheme (Table 2) cooked Frigate tuna was evaluated in parallel to the quality index method. High score (10) gives for the best quality and the low (3) score for the poorest quality. Scores for Frigate tuna stored 22 days in ice were found to be significantly different from those stored in ice for 1 to 7 days. The positive odour attributes, seaweed odour decreased with storage in ice and were hardly detectable after 12 days. The sour flavour decreased to unacceptable level of Frigate tuna for human consumption after 22 days in ice. Figure 3 shows the result of assessment of cooked fish using Torry scale during storage days in ice.




Figure 1. Relationship between Quality Index of Frigate tuna and storage period in ice.






Figure 2. Average scores of each quality attribute assessed with QIM scheme for Frigate tuna stored in ice against days in ice.

Quality index for Frigate tuna


Eyes: Colour of pupil




Gills: Colour




Gills: Smell



Gills: Mucus



Blood: Colour



Fillets: Colour




Figure 2. Average scores of each quality attribute assessed with QIM scheme for Frigate tuna stored in ice against days in ice (Continued).

Viscera




Figure 2. Average scores of each quality attribute assessed with QIM scheme for Frigate tuna stored in ice against days in ice (Continued).




Figure 3. Quality scores of cooked fish based on odour and flavour during storage in ice

Quality index for Frigate tuna




Figure 4. Hydrogen Sulphide (H_2S) producing bacteria (spoilage bacteria) and total viable counts (TVC) during storage in ice.

A similar trend was evident in flavour as for odour. Positive attributes such as seaweed, sweet, metallic, oily flavour decreased with storage days in ice, while the negative attributes rancid, sour flavours increased with storage days. Especially the sour flavour was dominant after 22 days of storage, indicating that the Frigate tuna was of marginal acceptability. Some panelists were unable to detect some of the attributes.

Microbial counts

Total viable counts (TVC) increased with storage time starting from 7 days of storage (Fig. 4). When the fish reached the limits of acceptability (sensory score of 4 in Torry scheme), TVC reached level of 10^6 cfu/g.

Counts of H_2S producing bacteria could be detected in fish after 16 days of storage. Figure 4 shows that they were absent at the beginning and at the end of shelf life the counts were reached above 10^2 cfu/g.

As shown in figure 5, QI scores and log TVC ($R^2 = 0.868$) are highly correlated.




Figure 5. Correlation between bacterial counts and quality index score.




Figure 6. Total volatile nitrogen (TVN) and trimethylamine (TMA) in raw Frigate tuna during the storage period.

Chemical analysis

During storage in ice the concentrations of the Total Volatile Nitrogen (TVN) in tuna increased with time while Tri Methyl Amine (TMA) content remained more or less constant (Fig. 6). At the rejection time, TVN has reached the concentration of around 35 mg/100g and TMA amount was 2.5mg/100g. The TVN level is highly correlated to QI score (Fig. 7).

Quality index for Frigate tuna




Figure 7. Correlation between TVN quality index score.

pH measurements

The pH value fluctuated slightly until 16 days, increased drastically until day 19 and decreased until 22 days (Fig. 8).

Photographs were used as guides for assessors to assess Frigate tuna by the QIM scheme on the basis of colour/appearance of whole fish, mucus and colour of gills and cornea, form and colour of pupil. The photographs of Frigate tuna after 1, and 22 days in ice are shown in Figure 9.




Figure 8. Changes in pH during storage of ungutted Frigate tuna in ice.

Frigate tuna stored 1 day in ice



Frigate tuna stored 22 days in ice



Figure 9: Photographs of Frigate tuna at three different stages, at the beginning of storage in ice (1 day in ice) and at rejection limit (22 days in ice).

Discussion

The increase in discoloration of skin of Frigate tuna during the period of storage in ice can perhaps be connected to how carotenoids are bound in the flesh. The carotenoid-binding proteins or lipoproteins are known to break down in the deterioration process of autolysis and bacterial activity (Torrissen *et al.* 1989).

The stiffness of fish was scored 0 at day 1 as the Frigate tuna was at rigor. The resolution of rigor causes the muscle to relax again and through storage in ice, the flesh becomes soft due to autolysis influenced by both fish muscle enzyme and microbial enzymes, (Nielsen 1995; Gill 1995).

The scores for the quality attributes of eyes increased constantly throughout the storage time in ice. At the end of shelf life cornea become concave. Nevertheless enough information on form and colour of pupil could not be obtained. At the beginning of the storage time when the Frigate tuna was very fresh, the odour of gills was described as fresh sea weedy or neutral. This is probably because newly caught fish contains low levels of

Quality index for Frigate tuna

volatile compounds (Olafsdottir and Fleurence 1997). The score for colour of blood was close to 0 until 7 days in ice. The blood in the abdomen was blood red at first, but later on it became brownish. When the fish meat exists in the form of oxymyoglobin, this compound is oxidized to meta myoglobin, which is brown in colour. The increase in these negative odour attributes is caused by activity of spoilage bacteria producing substances of bad smell and oxidation of fat (Gram 1995; Jorgensen 1995). Milo and Grosh (1996) analyzed various odourants in salmon of different freshness levels. They found propionadehyde and (Z)-1,5-octadien-3-one as the most potent high volatile odorants in cooked fresh salmon samples. The odour of these compounds is described as sweet and metallic (respectively). Various odourants detected from cooked salmon (Milo and Grosh 1996) may also be responsible for the flavour and odour of cooked Frigate tuna. The odour and flavour were attributed to acetaldehyde (sweet), hexanal and (Z, Z)-3,6-nonaldienal. The rancid odour increased in the last days of storage. The rancid flavour is known to be caused by formation of volatile oxidation products such as aldehydes and ketones (Milo and Grosh, 1996). At the beginning, microbial counts in flesh of Frigate tuna were very low. It implies that flesh of healthy, live or newly caught fish is sterile as the immune system of the fish prevents the bacteria from growing in the flesh. When fish dies, the immune system collapses and during storage, bacteria invade the flesh (Gram 1995). Similar results were noted by Lande and Rora (1999). Magnusson (1987) found TVC and H₂S producing bacteria in farmed salmon, whose the total viable count reached 10⁵ cfu/g in flesh after the storage time of 22 days. In ice stored Nile perch the total counts were at 10⁹ cfu/g on the day before the fish was rejected (Gram et al. 1989).

For the TVN concentration in muscle tissue of certain fish species a legal limit has been fixed by the EC-commission at 25mg/100g. TVN is produced mainly by H₂S producing bacteria. In the present study, that is detected after 16 days in ice. TMA is a pungent volatile amine often associated with the typical "fishy" odour of spoiling seafood. TMA and many other amines become volatile at elevated pH.

The low value of pH in first day of storage could be explained by "post mortem" glycolysis that results in the accumulation of lactic acid, which in turn lower the pH of muscle. Huss (1995) has reported pH drops from 6.8 to 6.1-6.5 in cod and 6.8 to 5.4-5.6 in tuna and halibut. The increase in pH with storage time is probably caused by formation of basic compounds (Huss 1988).

The changes occurring in the outer appearance of Frigate tuna with storage days in ice are presented in Figure 9. The appearance of the skin changes from being pearl-shiny to dull, the eyes change from being black and convex to grey and sunken. The gills change from red with clear mucus to

brownish with clotted, brown mucus. All those changes are listed in QIM scheme for Frigate tuna, with photographs when possible, making it easier to assess the freshness stage of Frigate tuna.

Acknowledgement

The authors wish to acknowledge with thanks the cooperation extended by Mrs. K.S. Hettiarachchi, Mrs. J.M. Chandrika and Mr. S.P. Jayasooriya during the course of the study. This project was funded by NARA.

References

- Bremner H.A. 1985.
A convenient easy to use system for estimating the quality of chilled seafood. *Fish Processing Bulletin* 7: 59-703.
- Bremner H.A., J. Olley & A.M.A. Vail 1985.
Estimating time-temperature effects by a rapid systematic sensory method. In: *Proceedings of an international symposium on seafood quality determination, Alaska, U.S.A.* (D.E. Kramer and L. Liston eds), Elsevier, Amsterdam. 413-435 pp.
- EU directive 1991.
Council Directive 91/493 /EEC.
- Gill T. 1995.
Autolytic changes. In: *Quality and quality changes in fresh fish.* (H.H. Huss ed.). FAO technical paper, No. 348. FAO, Rome. 45-50 pp.
- Gram L. 1995.
Bacteriological changes. In: *Quality and quality changes in fresh fish.* (H.H. Huss ed.). FAO technical paper, No. 348. FAO, Rome. 51-63 pp.
- Gram L., G. Trolle & H.H. Huss 1989.
Detection of specific spoilage bacteria from fish stored at low (0°C) and high (20°C) temperatures. *Journal of Food Microbiology* 4: 65-72.
- Gram L., J. Oundo & J. Bon 1989.
Storage life of Nile perch (*Lates niloticus*) dependent on storage temperature and initial bacterial load. *Tropical Science* 29: 221-236.
- Horwits W. 1980.
Official Methods of Analysis Association of Official Analytical Chemists, Washington. DC-20044, USA.

Quality index for Frigate tuna

- Huss H.H. 1988.
Fresh fish quality and quality changes. FAO Fisheries Series 29.
FAO, Rome. 27-61 pp.
- Jorgensen B. 1995. Lipid oxidation and hydrolysis. In: Quality and quality changes in fresh fish (H.H. Huss ed.). FAO technical paper No. 348.
FAO, Rome. 64-67 pp.
- Lande A. & M.B. Rora 1999.
Holdbarhet av iset laks. Norsk Fiskeoppdrett nr. 5a: 6-7.
- Magnusson H. 1987.
Geymslupól a eldislaxi. Rannsóknastofnum fiskidnadarins, Reykjavík.
- Milo C. & W.Grosch 1996.
Changes in the odorants of boiled Salmon and Cod as affected by the storage of the raw material. Journal of Agriculture and Food Chemistry. 44: 2366-2371.
- Nielsen J. 1995.
Sensory methods. In: Quality and quality changes in fresh fish (H.H. Huss ed.). FAO technical paper No. 348. FAO, Rome. 130-139 pp.
- Olafsdottir G. & L. Fleurence 1997.
Evaluation of fish freshness using volatile compounds-classification of volatile compounds in fish. In: Methods to determine the freshness of fish (G. Olafsdottir et al. eds). Proceedings of the final meeting of the concerted action. Evaluation of fish freshness. Nantes 55-69 pp.
- Torrissen O.J., R.W. Hardy & Shearer 1989. Pigmentation of salmonoids - carotenoid deposition and metabolism. CRC Crit. Rev. Aquatic Science 1: 209-225.
- The Gazette of the Democratic Socialist Republic of Sri Lanka. Extraordinary. Fish products (Export) Regulation 1998. No. 1230/14, 03.04.2002.